

DESCRIPTION

HERBICIDAL COMPOSITION HAVING THE HERBICIDAL EFFECT
ENHANCED, AND METHOD FOR ENHANCING THE HERBICIDAL EFFECT
TECHNICAL FIELD

5 The present invention relates to a herbicidal
composition containing a sulfonylurea compound or its
salt, as a herbicidally active ingredient, and having the
herbicidal effect enhanced.

BACKGROUND ART

10 Heretofore, various herbicides have been used for
the purpose of protecting crop plants, and it has been
common to incorporate an adjuvant capable of enhancing
their herbicidal effects for the purpose of e.g. reducing
the dosage of the herbicidally active ingredient.

15 GB Patent Publication No. 2309904 discloses a
mixture comprising a dicarboxylic acid type compound and
an ethoxylated and/or propoxylated triglyceride, and it
is disclosed that the herbicidal effect can be enhanced
if this mixture is combined with a herbicide. Further,
20 EP 0598515A discloses that the herbicidal effect of a
certain specific sulfonylurea compound can be improved by
employing an ethoxylated fatty amine type surfactant, and
a vegetable oil and/or a mineral oil. However, these
references disclose nothing about use of an alkoxylated
25 glyceride to enhance the herbicidal effect of a
herbicidal sulfonylurea compound or its salt.

A sulfonylurea herbicide which is commonly and

commercially available is one of herbicides having small application dosages. However, from the viewpoint of the environmental problem or economical efficiency, it is desired to further reduce its application dosage. On the other hand, it is possible to reduce the application dosage of a sulfonylurea herbicide by incorporating an adjuvant. However, if the application amount of the adjuvant itself increases, it may adversely affect the environment or the economical efficiency. Thus, it is desired to reduce the application dosages of both the sulfonylurea herbicide and the adjuvant.

Under these circumstances, the present inventors have conducted various studies with an aim to remarkably enhance the herbicidal effect of a sulfonylurea herbicide thereby to reduce its application dosage, and as a result, have accomplished the present invention.

DISCLOSURE OF THE INVENTION

Namely, the present invention provides a herbicidal composition comprising a herbicidal sulfonylurea compound or its salt, and an alkoxyated glyceride; a method for controlling undesired plants by means of such a herbicidal composition; and a method for enhancing the herbicidal effect of a sulfonylurea compound or its salt, by means of an alkoxyated glyceride.

In the present invention, the herbicidal effect of the sulfonylurea compound or its salt can be remarkably improved by the alkoxyated glyceride. Accordingly, it

is thereby possible to provide a herbicidal composition having the herbicidal effect remarkably improved, whereby the application dosage of the sulfonylurea compound or its salt can be reduced, and a method for remarkably enhancing the herbicidal effect of the sulfonylurea compound or its salt. Further, the present invention brings about a merit such that the types of plants against which the herbicidal effect extends will increase, or the period for application can be extended.

10 BEST MODE FOR CARRYING OUT THE INVENTION


The herbicidal composition of the present invention comprises a herbicidal sulfonylurea compound or its salt, and an alkoxyated glyceride. Further, the present invention provides a method for controlling undesired plants or inhibiting their growth, which comprises applying (a) a herbicidally effective amount of a herbicidal sulfonylurea compound or its salt, and (b) an effective amount of an alkoxyated glyceride, to the undesired plants or to a place where they grow. The present invention is carried out, for example, (1) by formulating a sulfonylurea compound or its salt by means of various additives, diluting the formulation together with an alkoxyated glyceride with e.g. water, and applying it to the undesired plants or to a place where they grow, or (2) by formulating a sulfonylurea compound or its salt and an alkoxyated glyceride together with various additives, diluting the formulation with e.g.

water or without diluting it, and applying it to the undesired plants or to a place where they grow. In the present invention, in the application to the undesired plants or to a place where they grow, the application to
5 the undesired plants themselves, e.g. the foliar application, is preferred.

In the present invention, the sulfonylurea compound or its salt may be used in combination with one or more other herbicidal compounds. Namely, the present
10 invention can be carried out by a method for controlling undesired plants or inhibiting their growth by such a combined use of other herbicidal compounds, which comprises applying (a) a herbicidally effective amount of a herbicidal sulfonylurea compound or its salt, (b) an
15 effective amount of an alkoxyated glyceride, and (c) a herbicidally effective amount of other herbicidal compound(s), to the undesired plants or to a place where they grow. For example, it may be carried out (1) formulating a sulfonylurea compound or its salt and other
20 herbicidal compound(s) separately or all together, by means of various additives, diluting the formulation together with an alkoxyated glyceride with e.g. water, and applying it to the undesired plants or to a place where they grow, (2) by formulating a sulfonylurea
25 compound or its salt, and an alkoxyated glyceride, together with various additives, diluting the formulation together with a separately formulated other herbicidal

compound(s) with e.g. water, and applying it to the undesired plants or to a place where they grow, (3) by formulating a sulfonylurea compound or its salt, other herbicidal compound(s) and an alkoxyated glyceride, together with various additives, diluting the formulation with e.g. water or without diluting it, and applying it to the undesired plants or to a place where they grow, or (4) by formulating other herbicidal compound(s) and an alkoxyated glyceride together with various additives, diluting the formulation together with a separately formulated sulfonylurea compound or its salt with e.g. water, and applying it to the undesired plants or to a place where they grow.

The sulfonylurea compound is a compound having the following partial structure:



(wherein X is a hydrogen atom or an alkyl group), and it may, for example, be amidosulfuron, azimsulfuron, bensulfuron-methyl, chlorimuron-ethyl, chlorsulfuron, cinosulfuron, cyclosulfamuron, ethametsulfuron-methyl, ethoxysulfuron, flazasulfuron, flucetosulfuron, flupyrsulfuron, foramsulfuron, halosulfuron-methyl, imazosulfuron, iodosulfuron, mesosulfuron-methyl, metsulfuron-methyl, nicosulfuron, oxasulfuron, primisulfuron-methyl, prosulfuron, pyrazosulfuron-ethyl, rimsulfuron, sulfometuron-methyl, sulfosulfuron,

thifensulfuron-methyl, triasulfuron, tribenuron-methyl, trifloxysulfuron, triflusulfuron-methyl, or tritosulfuron. Among them, flazasulfuron, foramsulfuron, nicosulfuron, rimsulfuron, trifloxysulfuron or
5 tritosulfuron is preferred. Among them, nicosulfuron is particularly preferred.

As the salt of such a sulfonylurea compound(s), various types may be mentioned. It may, for example, be a salt with an alkali metal such as sodium or potassium,
10 a salt with an alkaline earth metal such as magnesium or calcium, or a salt with an amine such as monomethylamine, dimethylamine or triethylamine.

The alkoxyated glyceride is one to enhance the herbicidal effect of the sulfonylurea compound or its
15 salt, and it may, for example, be as follows.


As the alkoxyated glyceride, a monoglyceride, a diglyceride and a triglyceride are present, and all of them are included in the present invention.

The alkoxyated glyceride has one or more alkylene
20 oxide moieties at optional positions of the glyceride. Such an alkylene oxide moiety may, for example, be ethylene oxide, propylene oxide, a copolymer thereof or a block copolymer thereof. The average addition molar amount of the alkylene oxide is from 1 to 200 mols,
25 preferably from 2 to 150 mols, more preferably from 10 to 100 mols.

The glyceride moiety of the alkoxyated glyceride

may, for example, be a fatty acid ester having glycerin or its derivative bonded with a fatty acid group or its derivative, or a phosphoglyceride having glycerin or its derivative bonded with a phosphoric acid group or its derivative. Particularly preferred is a fatty acid ester having glycerin or its derivative bonded with a fatty acid group or its derivative.

The above glycerin or its derivative may, for example, be a compound represented by the following formula (I):



(wherein R is a hydrogen atom or a C₁₋₆ alkyl group, and n is an integer of from 0 to 6.)

The above fatty acid group may be either saturated or unsaturated and may be linear or branched, and its carbon number may be from 4 to 24, preferably from 10 to 20. The fatty acid may, for example, be a saturated fatty acid such as butyric acid, n-caproic acid, caprylic acid, n-capric acid, lauric acid, myristic acid, palmitic acid, stearic acid, isostearic acid or arachic acid, or an unsaturated fatty acid such as palmitoleic acid, oleic acid, linoleic acid, linolenic acid, monoctic acid, arachidonic acid or docosaheanoic acid. Among them, palmitic acid, stearic acid, isostearic acid or oleic

acid is, for example, preferred.

The above fatty acid group may be cyclic and may contain one or more hetero atoms in such a cyclic group, or the methylene moiety may be converted to oxo or thioxo. As a specific example, a pyroglutamic acid group may be mentioned. Further, the fatty acid group may be substituted by a hydroxyl group at an optional position. As a specific example, a ricinoleic acid group may be mentioned.

Further, the alkoxyated glyceride may have an alkylene oxide at the terminal of the fatty acid group (on the side opposite to the bonding cite with glycerin), and at the terminal of such an alkylene oxide, a fatty acid group may further be bonded.

The above phosphoglyceride may, for example, be phosphatidylcholine, phosphatidylethanolamine, phosphatidylserine, phosphatidylinositol, phosphatidylglycerol or diphosphatidylglycerol.

The following may, for example, be mentioned as specific examples of the alkoxyated glyceride.

Polyoxyethylene hydrogenated castor oil (tradenames: EMANON CH-25, EMANON CH-80; EMALOX HC-5, EMALOX HC-100; Sorpol HC-10, Sorpol HC-20, Sorpol HC-40, Sorpol HC-50, Sorpol HC-80, Sorpol HC-100, Sorpol HC-150; PEGNOL HC-30; NIKKOL HCO-5, NIKKOL HCO-10, NIKKOL HCO-20, NIKKOL HCO-30, NIKKOL HCO-40, NIKKOL HCO-50, NIKKOL HCO-60, NIKKOL HCO-80, NIKKO HCO-100, etc.)

Polyoxyethyleneglyceryl monostearate (tradenames:
EMALEX GM-5, EMALEX GM-40, etc.)

Polyoxyethylene castor oil (tradenames: EMALEX C-20,
EMALEX C-50, etc.)

5 Polyoxyethylene glyceryl triisostearate (tradenames:
EMALEX GWIS-303, EMALEX GWIS-340, EMALEX GWIS-360, EMALEX
GWIS-340EX, etc.)

Polyoxyethylene glyceryl monoisostearate
(tradenames; EMALEX GWIS-103, EMALEX GWIS-115, EMALEX
10 GWIS-125, EMALEX GWIS-160, EMALEX GWIS-160EX, etc.)

Polyoxyethylene glyceryl tristearate (tradenames:
EMALEX GWS-303, EMALEX GWS-320, etc.)

Polyoxyethylene glyceryl distearate (tradenames:
EMALEX GWS-204, etc.)

15 Polyoxyethylene glyceryl trioleate (tradename:
EMALEX GWO-303, EMALEX GWO-360, etc.)

Polyoxyethylene hydrogenated castor oil
monoisostearate (tradenames: EMALEX RWIS-105, EMALEX
RWIS-158, EMALEX RWIS-150EX, etc.)

20 Polyoxyethylene hydrogenated castor oil
triisostearate (tradenames: EMALEX RWIS-305, EMALEX RWIS-
360, EMALEX RWIS-360EX, etc.)

Polyoxyethylene hydrogenated castor oil monolaurate
(tradenames: EMALEX RWL-120, EMALEX RWL-160, etc.)

25 Polyoxyethylene 1,1,1-trimethylolpropane tristearate
(tradenames: EMALEX TPS-303, EMALEX TPS-310, etc.)

Polyoxyethylene 1,1,1-trimethylolpropane

trimyrystate (tradenames: EMALEx TPM-303, EMALEx TPM-330, etc.)

Polyoxyethylene 1,1,1-trimethylolpropane distearate (tradenames: EMALEx TPS-203, EMALEx TPS-205, etc.)

5 Polyoxyethylene 1,1,1-trimethylolpropane triisostearate (tradenames: EMALEx TPIS-303, EMALEx TPIS-350, etc.)

Polyoxyethylene hydrogenated castor oil pyroglutamic acid isostearate (tradenames: PYROTER CPI-30, PYROTER
10 CPI-60, etc.)

Polyoxyethylene glyceryl pyroglutamic acid isostearate (tradenames: PYROTER GPI-25, etc.)

Among those represented by the above tradenames, EMANONS are manufactured by Kao Corporation, EMALEx and
15 PYROTER are manufactured by Nihon Emulsion Co., Ltd., Sorpol and PEGNOL are manufactured by TOHO Chemical Industry Co., Ltd., and NIKKOLS are manufactured by NIKKO CHEMICALS CO., LTD.

Among the above alkoxyated glycerides,
20 polyoxyethylene hydrogenated castor oil, polyoxyethylene glyceryl triisostearate, polyoxyethylene glyceryl monoisostearate, polyoxyethylene 1,1,1-trimethylolpropane triisostearate, polyoxyethylene hydrogenated castor oil pyroglutamic acid isostearate, or polyoxyethylene
25 glyceryl pyroglutamic acid isostearate may, for example, be preferred.

As other herbicidal compound(s) which can be used in

combination with the sulfonylurea compound or its salt, the compound groups of the following (1) to (10) (common names including those which are applied for ISO) may, for example, be mentioned. Even when not specifically
5 mentioned, in a case where such compounds have salts, alkyl esters or various structural isomers such as optical isomers, they are, of course, all included.

(1) Those which are believed to exhibit herbicidal effects by disturbing hormone activities of plants, such
10 as a phenoxy type such as 2,4-D, 2,4-DP, MCPA, MCPB, MCPP or naproanilide, an aromatic carboxylic acid type such as 2,3,6-TBA, dicamba, dichlobenil, picloram, triclopyr or clopyralid, and others such as benazolin, quinclorac, quinmerac, diflufenzopyr and thiazopyr.

15 (2) Those which are believed to exhibit herbicidal effects by inhibiting photosynthesis of plants, such as, a urea type such as chlorotoluron, diuron, fluometuron, linuron, isoproturon, metobenzuron or tebuthiuron, a triazine type such as simazine, atrazine, atratone,
20 simetryn, prometryn, dimethametryn, hexazinone, metribuzin, terbuthylazine, cyanazine, ametryn, cybutryne, triaziflam or propazine, a uracil type such as bromacil, lenacil or terbacil, an anilide type such as propanil or cypromid, a carbamate type such as swep,
25 desmedipham or phenmedipham, a hydroxybenzonitrile type such as bromoxynil, bromoxynil-octanoate or ioxynil, and others such as pyridate, bentazon and amicarbazone.

(3) Quaternary ammonium salt type such as paraquat or diquat, which is believed to be converted to free radicals by itself to form active oxygen in the plant body.

5 (4) Those which are believed to exhibit herbicidal effects by inhibiting chlorophyll biosynthesis of plants and abnormally accumulating a photosensitizing peroxide substance in the plant body, such as a diphenylether type such as nitrofen, chlomethoxyfen, bifenox, acifluorfen-
10 sodium, fomesafen, oxyfluorfen, lactofen or ethoxyfen-ethyl, a cyclic imide type such as chlorphthalim, flumioxazin, flumiclorac-pentyl or fluthiacet-methyl, and others such as oxadiargyl, oxadiazon, sulfentrazone, carfentrazone-ethyl, thidiazimin, pentoxazone,
15 azafenidin, pyraflufen-ethyl, benzfendizone, butafenacil, metobenzuron, cinidon-ethyl, flupoxam, fluazolate, profluazol and pyrachlonil.

 (5) Those which are believed to exhibit herbicidal effects characterized by whitening activities by
20 inhibiting chromogenesis of plants such as carotenoids, such as a pyridazinone type such as norflurazon or metflurazon, a pyrazole type such as pyrazolate, pyrazoxyfen or benzofenap, and others such as amitrol, fluridone, flurtamone, diflufenican, methoxyphenone,
25 clomazone, sulcotrione, mesotrione, isoxaflutole, difenzoquat, isoxachlortole, benzobicyclone, picolinofen and beflubutamid.

(6) Those which exhibit strong herbicidal effects specifically to gramineous plants, such as an aryloxyphenoxypropionic acid type such as diclofop-methyl, flamprop-M-methyl, pyriphenop-sodium, fluazifop-
5 butyl, haloxyfop-methyl, quizalofop-ethyl, cyhalofop-butyl or fenoxaprop-ethyl, and a cyclohexanedione type such as alloxydim-sodium, clethodim, sethoxydim, tralkoxydim, butroxydim, tepraloxydim, caloxydim or clefoxydim.

10 (7) Those which are believed to exhibit herbicidal effects by inhibiting an amino acid biosynthesis of plants, such as a triazolopyrimidinesulfonamide type such as flumetsulam, metosulam, diclosulam, cloransulam-methyl, florasulam, metosulfam or penoxsulam, an
15 imidazolinone type such as imazapyr, imazethapyr, imazaquin, imazamox, imazameth, imazamethabenz or imazapic, a pyrimidinylsalicylic acid type such as pyrithiobac-sodium, bispyribac-sodium, pyriminobac-methyl, pyribenzoxim or pyriftalid, a
20 sulfonylaminocarbonyltriazolinone type such as flucarbazone or procarbazon-sodium, and others such as glyphosate-ammonium, glyphosate-isopropylamine, glufosinate-ammonium and bialaphos.

(8) Those which are believed to exhibit herbicidal
25 effects by inhibiting cell mitoses of plants, such as a dinitroaniline type such as trifluralin, oryzalin, nitralin, pendimethalin or ethalfluralin, an organic

phosphorus type such as amiprofos-methyl, butamifos, anilofos or piperophos, a phenylcarbamate type such as chlorpropham or barban, a cumylamine type such as daimuron, cumyluron or bromobutide, and others such as
5 asulam, dithiopyr and thiazopyr.

(9) Those which are believed to exhibit herbicidal effects by inhibiting protein biosynthesis or lipid biosynthesis of plants, such as, a thiocarbamate type such as EPTC, butylate, molinate, dimepiperate,
10 fluazolate, esprocarb, thiobencarb, pyributicarb or trialate, a chloroacetamide type such as alachlor, butachlor, pretilachlor, metolachlor, S-metolachlor, thenylchlor, pethoxamid, dimethenamid, acetochlor or propachlor, and others such as etobenzanid, mefenacet,
15 flufenacet, tridiphane, cafenstrole, fentrazamide, oxaziclomefone and indanofan.

(10) Those which are believed to exhibit herbicidal effects by being parasitic on plants, such as Xanthomonas campestris, Epicoccosurus nematosurus, Exserohilum
20 monoseras and Drechsrela monoceras.

In the present invention, a coadjuvant may be employed, as the case requires for the purpose of more distinctly enhancing the herbicidal effects of the sulfonylurea compound or its salt, increasing the types
25 of plants against which the herbicidal effects are effective, or expanding the application period.

As such a coadjuvant, a chelating agent, a nitrogen-

containing fertilizer or a cationic surfactant may, for example, be mentioned. Among them, a chelating agent or a nitrogen-containing fertilizer is preferred. When coadjuvants are to be applied, two or more of them may
5 suitably be used in combination, as the case requires.

Specific examples of the chelating agent include aminopolycarboxylic acids and aliphatic carboxylic acids. The aminopolycarboxylic acids may, for example, be ethylenediaminetetraacetic acid (EDTA), iminodiacetic
10 acid (IDA), nitrilotriacetic acid (NTA), ethylene glycol bis(2-aminoethyl ether)-N,N,N',N'-tetraacetic acid (EGTA), diethylenetriamine-N,N,N',N',N'-pentaacetic acid (DTPA), cyclohexanediaminetetraacetic acid (CDTA), or their salts. Further, the aliphatic carboxylic acids
15 may, for example, be citric acid, malic acid, oxalic acid, lactic acid, gluconic acid, heptonic acid, or their salts. Two or more of them may suitably be used in combination, as the case requires.

Specific examples of the nitrogen-containing
20 fertilizer include ammonium salts such as ammonium sulfate, ammonium chloride, ammonium phosphite and ammonium phosphate; and nitrates such as ammonium nitrate, potassium nitrate and calcium nitrate. Two or more of them may suitably be used in combination, as the
25 case requires.

Specific examples of the cationic surfactant include tertiary amines such as alkoxylated fatty acid amines;

salts of tertiary amines; and quaternary ammonium salts. Two or more of them may suitably be used in combination, as the case requires.

As the above-mentioned salts of amino polycarboxylic acids or aliphatic carboxylic acids, various salts may be mentioned. For example, a salt with an alkali metal such as sodium or potassium, a salt with an alkaline earth metal such as magnesium or calcium, or a salt with an amine such as monomethylamine, dimethylamine or triethylamine, may be mentioned.

In the present invention, the blend ratio of the sulfonylurea compound or its salt to the alkoxyated glyceride cannot generally be defined, since it may be suitably changed depending upon the type of the sulfonylurea compound or the alkoxyated glyceride, the formulation, the climate condition, the type or size of the plants to be controlled, etc. However, it may, for example, be within a range of from 16:1 to 1:6000, more preferably from 8:1 to 1:1000, further preferably from 2:1 to 1:600, still further preferably from 1:2 to 1:50, by weight ratio.

In a case where a coadjuvant is used in the present invention, the blend ratio cannot generally be defined, since it may be suitably changed depending upon the type of the sulfonylurea compound, the type of the alkoxyated glyceride, the type of the coadjuvant, the formulation, the climate conditions, the type and size of the plants

to be controlled, etc. However, the blend ratio of the alkoxyated glyceride and the coadjuvant may, for example, be within a range of from 4000:1 to 1:5, preferably from 500:1 to 2:1, by weight ratio.

5 Further, if the blend ratio is represented for each mode of practical applications, the following may, for example, be mentioned. (1) In a case where the sulfonylurea compound or its salt is formulated by means of various additives, then diluted together with the
10 alkoxyated glyceride with e.g. water and applied to undesired plants or to a place where they grow, and the formulated sulfonylurea compound or its salt and alkoxyated glyceride are diluted with from 10 to 3000 liters/ha, preferably from 50 to 2000 liters/ha, more
15 preferably from 100 to 1000 liters/ha, of water, the alkoxyated glyceride is added in an amount of from 0.005 to 4 wt%, preferably from 0.01 to 2 wt%, more preferably from 0.02 to 0.5 wt%, based on the diluted liquid. (2)
In a case where the sulfonylurea compound or its salt,
20 and the alkoxyated glyceride, are formulated together with various additives, then diluted with e.g. water or without dilution, and applied to undesired plants or to a place where they grow, the sulfonylurea compound or its salt, and the alkoxyated glyceride, may be blended so
25 that their ratio will be in the above-mentioned weight ratio range. (3) In a case where one or more other herbicidal compounds are used in combination with the

sulfonylurea compound or its salt, such can be carried out in accordance with the blend ratio in the above case (1) or (2). (4) In a case where a coadjuvant is further employed and they are diluted with water in the same manner as described above, in each of the above cases (1), (2) and (3), it may be added in an amount of from 0.0001 to 0.05 wt%, preferably from 0.001 to 0.01 wt%, based on the diluted liquid.

In the present invention, (1) in a case where the sulfonylurea compound or its salt, and other herbicidal compound(s), are, separately or together, formulated by means of various additives, (2) in a case where the sulfonylurea compound or its salt, and the alkoxyated glyceride, are formulated together with various additives, (3) in a case where the sulfonylurea compound or its salt, other herbicidal compound(s) and the alkoxyated glyceride, are formulated together with various additives, or (4) in a case where the sulfonylurea compound or its salt, other herbicidal compound(s), the alkoxyated glyceride and the coadjuvant, are formulated together with various additives, they may be formulated into various formulations. For example, various formulations such as a wettable powder, water-dispersible granules, a water-based suspension concentrate, an oil-based suspension concentrate, a gel formulation, an emulsifiable concentrate, water soluble granules, an emulsion, a

microemulsion, a suspoemulsion, and a multiple emulsion, may be mentioned. Various additives which can be used here may be any additives so long as they are commonly used in this technical field, and for example, a

5 surfactant, a carrier, a solvent, a vegetable oil, a mineral oil, an anti-settling agent, a thickener, an antifoaming agent, an antifreezing agent, an antioxidant, an oil absorbent, a gelling agent, a filler, a dispersion stabilizer, a phytotoxicity reducing agent, an anti-mold

10 agent, a binder, a stabilizer, a disintegrator, a preservative, and an inorganic ammonium salt, may be mentioned. The following may, for example, be mentioned as specific examples of such various additives. Further, such formulations may be prepared in accordance with

15 methods commonly employed in this technical field.

The surfactant includes, for example, anionic surfactants such as a salt of fatty acid, a benzoate, an alkylsulfosuccinate, a dialkylsulfosuccinate, a polycarboxylate, a salt of alkylsulfuric acid ester, an

20 alkyl sulfate, an alkylaryl sulfate, an alkyl diglycol ether sulfate, a salt of alcohol sulfuric acid ester, an alkyl sulfonate, an alkylaryl sulfonate, an aryl sulfonate, a lignin sulfonate, an alkyl diphenyl ether disulfonate, a polystyrene sulfonate, a salt of

25 alkylphosphoric acid ester, an alkylaryl phosphate, a styrylaryl phosphate, a salt of polyoxyethylene alkyl ether sulfuric acid ester, a polyoxyethylene alkylaryl

ether sulfate, a polyoxyethylene styrylaryl ether sulfate, an ammonium polyoxyethylene styrylaryl ether sulfate, a salt of polyoxyethylene alkylaryl ether sulfuric acid ester, a polyoxyethylene alkyl ether phosphate, a salt of polyoxyethylene alkylaryl phosphoric acid ester, a polyoxyethylene styrylaryl ether phosphoric acid ester or its salt, a salt of a condensate of naphthalene sulfonate with formalin and a salt of a condensate of alkyl naphthalene sulfate with formalin;

10 nonionic surfactants such as a sorbitan fatty acid ester, a glycerin fatty acid ester, a fatty acid polyglyceride, a fatty acid alcohol polyglycol ether, acetylene glycol, acetylene alcohol, an oxyalkylene block polymer, a polyoxyethylene alkyl ether, a polyoxyethylene alkylaryl ether, a polyoxyethylene styrylaryl ether, a

15 polyoxyethylene glycol alkyl ether, a polyoxyethylene fatty acid ester, a polyoxyethylene sorbitan fatty acid ester, a polyoxyethylene glycerin fatty acid ester, a polyoxyethylene hydrogenated castor oil, and a

20 polyoxypropylene fatty acid ester, and cationic surfactants such as an alkoxylated fatty acid amine. If desired, two or more of them may suitably be used in combination.

The carrier or the filler may, for example, be

25 diatomaceous earth, slaked lime, calcium carbonate, talc, white carbon, kaoline, bentonite, a mixture of kaolinite and sericite, clay, sodium carbonate, sodium bicarbonate,

mirabilite, zeolite, starch, sodium chloride, ammonium phosphate, ammonium sulfate, ammonium chloride, sugar, urea, lactose, or glucose. If desired, two or more of them may suitably be used in combination.

5 The solvent may, for example, be water, solvent naphtha, paraffin, dioxane, acetone, isophorone, methyl isobutyl ketone, chlorobenzene, cyclohexane, dimethyl sulfoxide, dimethyl formamide, N-methyl-2-pyrrolidone, an alcohol, acetic acid, butyric acid, isopropyl acetate,
10 butyl acetate, benzene, an alkylbenzene or an alkylnaphthalene. If desired, two or more of them may suitably be used in combination.

 The vegetable oil may, for example, be olive oil, kapok oil, castor oil, papaya oil, camellia oil, coconut
15 oil, sesame oil, corn oil, rice bran oil, peanut oil, cottonseed oil, soybean oil, rapeseed oil, linseed oil, tung oil, sunflower oil, safflower oil, or a fatty acid derived from each of the above-mentioned oils or an alkyl ester of such a fatty acid. The mineral oil may, for
20 example, be an aliphatic hydrocarbon such as liquid paraffin or paraffinic petroleum, or an aromatic hydrocarbon such as an alkyl benzene or an alkyl naphthalene. If desired two or more of them may suitably be used in combination. The above fatty acid may, for
25 example, be a C₁₂₋₂₂ saturated or unsaturated fatty acid, such as lauric acid, palmitic acid, stearic acid, oleic acid, linolic acid, linoleic acid, erucic acid or

brassidic acid, and the alkyl ester thereof may, for example, a C₁₋₁₈ linear or branched alkyl ester, such as a methyl ester, a butyl ester, an isobutyl ester or an oleyl ester.

5 The anti-settling agent may, for example, be silica, an organic bentonite (bentonite-alkylamino complex), bentonite, white carbon or aluminum magnesium silicic acid. If desired, two or more of them may be suitably used in combination.

10 The thickener may, for example, be a heteropolysaccharide such as xanthan gum or guar gum, a water-soluble polymer such as polyvinyl alcohol, a sodium salt of carboxymethyl cellulose or sodium alginate, bentonite or white carbon. If desired, two or more of
15 them may suitably be used in combination.

 The antifoaming agent may, for example, be polydimethylsiloxane or acetylene alcohol. If desired, two or more of them may suitably be used in combination.

 The antifreezing agent may, for example, be ethylene
20 glycol, propylene glycol, glycerin or urea. If desired, two or more of them may suitably be used in combination.

 The oil-absorber may, for example, be silicon dioxide, hydrolyzed starch, kaoline, clay, talc, diatomaceous earth, a synthesized product of diatomaceous
25 earth/lime, asbestos, a mixture of kaolinite and sericite, calcium silicate, calcium carbonate, calcium carbonate silicate, acid clay, carbon black, graphite, a

pearlite processed product, alumina, titanium dioxide, basic magnesium carbonate, magnesium silicate aluminate, silica-alumina filler or magnesium silicate hydrate. If desired, two or more of them may suitably be used in
5 combination.

The gelling agent may, for example, be silica, organic attapulgite, clay, hydrogenated castor oil, a higher fatty acid ester, a higher alcohol, a salt of dialkyl sulfosuccinic acid ester, a benzoate, an alkyl
10 sulfate, a poly acrylic polymer, or a mixture of a poly acrylic acid co-polymer and water, or 12-hydroxystearic acid. If desired, two or more of them may suitably be used in combination.

The binder may, for example, be lignin sulfonic
15 acid, xanthan gum, carboxymethylcellulose or starch. If desired, two or more of them may suitably be used in combination.

The stabilizer may, for example, be urea.

The disintegrator may, for example, be a calcium
20 salt of carboxymethylcellulose, an inorganic salt such as ammonium sulfate, potassium chloride or magnesium chloride, and one showing a disintegrating activity among the above-mentioned surfactants, such as sodium lauryl sulfate, sodium dodecylbenzene sulfonate, or an ammonium
25 polyacrylate. If desired, two or more of them may suitably be used in combination.

The preservative may, for example, be formalin, p-

chloro m-xylenol or 1,2-benzisothiazolin-3-one. If desired, two or more of them may suitably be used in combination.

In the above-mentioned various formulations, the
5 blend proportions of various components cannot generally be defined, since they may suitably be changed depending upon the types of the blend components, the formulations or the application sites. However, a formulation may, for example, be prepared by incorporating the
10 sulfonylurea compound or its salt in a proportion of from 0.1 to 95 parts by weight, preferably from 2 to 85 parts by weight and incorporating, as the rest, various additives in a proportion of from 5 to 99.9 parts by weight, preferably from 15 to 98 parts by weight.
15 Further, in a case where the alkoxyated glyceride is incorporated in a proportion of from 0.1 to 94.9 parts by weight, preferably from 5 to 60 parts by weight, as the case requires, where other herbicidal compound(s) are incorporated in a proportion of from 0.1 to 94.9 parts by
20 weight, preferably from 0.5 to 75 parts by weight, as the case requires, or where the coadjuvant is incorporated in a proportion of from 0.1 to 94.9 parts by weight, preferably from 0.2 to 60 parts by weight, as the case requires, a formulation may be prepared by incorporating
25 various additives as the rest, so that the total would be 100 parts by weight. Now, the blend proportions of the respective components in some formulations will be .

exemplified below, but it should be understood that the present invention is by no means restricted to such specific formulations.

In the case of a wettable powder, a formulation may
5 be prepared by incorporating the sulfonylurea compound or its salt in a proportion of from 0.1 to 95 parts by weight, preferably from 5 to 85 parts by weight, blending a surfactant in a proportion of from 0.5 to 40 parts by weight, preferably from 5 to 30 parts by weight, and
10 blending as the rest, the carrier or filler in a proportion of from 4.5 to 99.4 parts by weight, preferably from 10 to 90 parts by weight. Further, in a case where the alkoxyated glyceride is incorporated in a proportion of from 0.1 to 94.9 parts by weight,
15 preferably from 10 to 60 parts by weight, as the case requires, where other herbicidal compound(s) are incorporated in a proportion of from 0.1 to 94.9 parts by weight, preferably from 0.5 to 75 parts by weight, as the case requires, where a coadjuvant is incorporated in a
20 proportion of from 0.1 to 94.9 parts by weight, preferably from 0.2 to 60 parts by weight, as the case requires, or where an oil absorbent is incorporated in a proportion of from 1 to 90 parts by weight, preferably from 1 to 50 parts by weight, as the case requires, a
25 formulation may be prepared by incorporating a carrier or filler as the rest, so that the total would be 100 parts by weight.

In the case of water-dispersible granules, a formulation may be prepared by incorporating the sulfonylurea compound or its salt in a proportion of from 0.1 to 95 parts by weight, preferably from 5 to 85 parts by weight, and the surfactant in a proportion of from 0.5 to 40 parts by weight, preferably from 5 to 30 parts by weight, and incorporating, as the rest, a carrier or filler in a proportion of from 4.5 to 99.4 parts by weight, preferably from 10 to 90 parts by weight.

Further, in a case where the alkoxylated glyceride is incorporated in a proportion of from 0.1 to 94.9 parts by weight, preferably from 10 to 60 parts by weight, as the case requires, where other herbicidal compound(s) are incorporated in a proportion of from 0.1 to 94.9 parts by weight, preferably from 0.5 to 75 parts by weight, as the case requires, where the coadjuvant is incorporated in a proportion of from 0.1 to 94.9 parts by weight, preferably from 0.2 to 60 parts by weight, as the case requires, where a binder is incorporated in a proportion of from 0.1 to 10 parts by weight, preferably from 0.5 to 5 parts by weight, as the case requires, where a disintegrator is incorporated in a proportion of from 0.1 to 40 parts by weight, preferably from 0.5 to 20 parts by weight, as the case requires, or where an oil absorbent is incorporated in a proportion of from 1 to 90 parts by weight, preferably from 1 to 50 parts by weight, as the case requires, a formulation may be prepared by

incorporating, as the rest, a carrier or filler, so that the total would be 100 parts by weight.

In the case of a water-based suspension concentrate, a formulation may be prepared by incorporating the

5 sulfonylurea compound or its salt in a proportion of from 0.1 to 60 parts by weight, preferably from 2 to 50 parts by weight, and a surfactant in a proportion of from 0.5 to 20 parts by weight, preferably from 1 to 15 parts by weight, and incorporating, as the rest, water in a

10 proportion of from 25 to 99.4 parts by weight, preferably from 30 to 97 parts by weight. Further, in a case where the alkoxyated glyceride is incorporated in a proportion of from 0.1 to 60 parts by weight, preferably from 5 to 40 parts by weight, as the case requires, where other

15 herbicidal compound(s) are incorporated in a proportion of from 0.1 to 60 parts by weight, preferably from 0.5 to 30 parts by weight, as the case requires, where a coadjuvant is incorporated in a proportion of from 0.1 to 60 parts by weight, preferably from 0.2 to 40 parts by

20 weight, as the case requires, where an antifoaming agent is incorporated in a proportion of from 0.05 to 3 parts by weight, preferably from 0.1 to 1 parts by weight, as the case requires, where an anti-freezing agent is incorporated in a proportion of from 0.5 to 10 parts by

25 weight, preferably from 2 to 10 parts by weight, as the case requires, where an anti-settling agent is incorporated in a proportion of from 0.1 to 5 parts by

weight, preferably from 0.5 to 3 parts by weight, as the case requires, where a thickener is incorporated in a proportion of from 0.1 to 5 parts by weight, preferably from 0.1 to 2 parts by weight, as the case requires, or
5 where a preservative is incorporated in a proportion of from 0.01 to 1 part by weight, preferably from 0.05 to 0.2 part by weight, as the case requires, a formulation may be prepared by incorporating water as the rest, so that the total would be 100 parts by weight.

10 In the case of an oil-based suspension concentrate, a formulation may be prepared by incorporating the sulfonylurea compound or its salt in a proportion of from 0.1 to 40 parts by weight, preferably from 2 to 35 parts by weight, and a surfactant in a proportion of from 1 to
15 30 parts by weight, preferably from 1 to 25 parts by weight, and incorporating, as the rest, a vegetable or mineral oil in a proportion of from 10 to 98.9 parts by weight, preferably from 20 to 97 parts by weight.
Further, in a case where the alkoxyated glyceride is
20 incorporated in a proportion of from 0.1 to 80 parts by weight, preferably from 5 to 60 parts by weight, as the case requires, where other herbicidal compound(s) are incorporated in a proportion of from 0.1 to 40 parts by weight, preferably from 0.5 to 30 parts by weight, as the
25 case requires, where a coadjuvant is incorporated in a proportion of from 0.1 to 40 parts by weight, preferably from 0.2 to 35 parts by weight, as the case requires,

where an anti-settling agent is incorporated in a proportion of from 0.1 to 5 parts by weight, preferably from 0.5 to 3 parts by weight, as the case requires, or where a stabilizer is incorporated in a proportion of
5 from 0.1 to 20 parts by weight, preferably from 1 to 10 parts by weight, as the case requires, a formulation may be prepared by incorporating a vegetable oil or mineral oil as the rest, so that the total would be 100 parts by weight.

10 In the case of a gel formulation, a formulation may be prepared by incorporating the sulfonylurea compound or its salt in a proportion of from 0.1 to 40 parts by weight, preferably from 2 to 30 parts by weight, a surfactant in a proportion of from 1 to 30 parts by
15 weight, preferably from 1 to 15 parts by weight, and a gelling agent in a proportion of from 0.1 to 50 parts by weight, preferably from 5 to 40 parts by weight, and incorporating, as the rest, a vegetable oil or mineral oil in a proportion of from 10 to 98.8 parts by weight,
20 preferably from 20 to 92 parts by weight. Further, in a case where the alkoxylated glyceride is incorporated in a proportion of from 0.1 to 60 parts by weight, preferably from 5 to 25 parts by weight, as the case requires, where other herbicidal compound(s) are incorporated in a
25 proportion of from 0.1 to 30 parts by weight, preferably from 0.5 to 20 parts by weight, as the case requires, where a coadjuvant is incorporated in a proportion of

from 0.1 to 40 parts by weight, preferably from 0.2 to 30 parts by weight, as the case requires, or where an anti-settling agent is incorporated in a proportion of from 0.1 to 3 parts by weight, preferably from 0.5 to 2 parts by weight, as the case requires, a formulation may be prepared by incorporating a vegetable oil or mineral oil as the rest, so that the total would be 100 parts by weight.

The herbicidal composition of the present invention is capable of controlling a wide range of weeds including, for example, sedge (or Cyperaceae) such as rice flatsedge (Cyperus iria L.) or purple nutsedge (Cyperus rotundus L.), grasses (or gramineae) such as barnyardgrass (Echinochloa crus-galli L.), crabgrass (Digitaria sanguinalis L.), green foxtail (Setaria viridis L.), goosegrass (Eleusine indica L.), wild oat (Avena fatua L.), johnsongrass (Sorghum halepense L.) or quackgrass (Agropyron repens L.), and broadleaves such as velvetleaf (Abutilon theophrasti MEDIC.), tall morning glory (Ipomoea purpurea L.), common lambsquarters (Chenopodium album L.), prickly sida (Sida spinosa L.), common purslane (Portulaca oleracea L.), redroot pigweed (Amaranthus retroflexus L.), sicklepod (Cassia obtusifolia L.), black nightshade (Solanum nigrum L.), pale smartweed (Polygonum lapathifolium L.), common chickweed (Stellaria media L.), common cocklebur (Xanthium strumarium L.), flexuous bittercress (Cardamine

flexuosa WITH.), henbit (Lamium amplexicaule L.) or three seeded copperleaf (Acalypha austalis L.), or inhibiting their growth, by applying it to such undesired plants or to a place where they grow, for example, by foliar application. Accordingly, its application range extends not only to crop plant fields but also agricultural fields such as orchards or mulberry fields and non-agricultural fields such as forest land, farm roads, play grounds, factory sites or grass plots. The sulfonylurea compound or its salt may be applied in an amount of 1 to 500 g/ha, preferably from 2 to 250 g/ha. Particularly, a herbicidal composition comprising nicosulfuron or its salt, and an alkoxylated glyceride, is capable of controlling noxious weeds or inhibiting their growth without presenting phytotoxicity to corn, and it is very useful as a herbicidal composition for corn fields. Nicosulfuron may be applied in an amount of from 2 to 400 g/ha, preferably from 5 to 200 g/ha. Further, it may be used in combination or admixture with other agricultural chemicals, fertilizers, phytotoxicity-reducing agents, etc., whereby it is expected to exhibit better effects or activities.

Some preferred embodiments of the present invention will be exemplified. However, the present invention is by no means restricted thereto.

(1) The above herbicidal composition comprising a) a herbicidally effective amount of the sulfonylurea

compound or its salt, and b) the alkoxyated glyceride in an effective amount to enhance the herbicidal activity.

(2) The above herbicidal composition comprising a) the sulfonylurea compound or its salt, b) the alkoxyated glyceride in an effective amount to enhance the herbicidal activity, and c) the coadjuvant in an effective amount to enhance the herbicidal activity.

(3) The above herbicidal composition characterized in that the sulfonylurea compound or its salt, and the alkoxyated glyceride, are formulated together with additives.

(4) The above herbicidal composition characterized in that the sulfonylurea compound or its salt, the alkoxyated glyceride and the coadjuvant are formulated together with additives.

(5) The above herbicidal composition characterized in that it comprises the sulfonylurea compound or its salt, and the alkoxyated glyceride, and it is in the form of a water-diluted liquid which can be applied to undesired plants or to a place where they grow.

(6) The above herbicidal composition characterized in that it comprises the sulfonylurea compound or its salt, the alkoxyated glyceride and the coadjuvant, and it is in the form of a water-diluted liquid which can be applied to undesired plants or to a place where they grow.

(7) The above herbicidal composition which further

contains a herbicidally effective amount of other herbicidal compound(s).

(8) The above herbicidal composition characterized in that the herbicidal effect of the sulfonylurea compound or its salt is enhanced by the alkoxy-
5 glyceride.

(9) The above herbicidal composition characterized in that the herbicidal effect of the sulfonylurea compound or its salt, is enhanced by the alkoxy-
10 glyceride and the coadjuvant.

(10) The above herbicidal composition characterized in that the herbicidal effect of the herbicidal composition comprising the sulfonylurea compound or its salt and other herbicidal compound(s), is enhanced by the
15 alkoxy- glyceride.

(11) The above herbicidal composition characterized in that the herbicidal effect of the herbicidal composition comprising the sulfonylurea compound or its salt, and other herbicidal compound(s), is enhanced by
20 the alkoxy- glyceride and the coadjuvant.

(12) The above-mentioned method for controlling undesired plants or inhibiting their growth, which comprises applying a) a herbicidally effective amount of the sulfonylurea compound or its salt, and b) the
25 alkoxy- glyceride in an effective amount to enhance the herbicidal activity, to undesired plants or to a place where they grow.

(13) The above method for controlling undesired plants or inhibiting their growth, which comprises applying a) a herbicidally effective amount of the sulfonylurea compound or its salt, b) the alkoxyated glyceride in an effective amount to enhance the herbicidal activity, and c) the coadjuvant in an effective amount to enhance the herbicidal activity, to undesired plants or to a place where they grow.

(14) The above method for controlling undesired plants or inhibiting their growth, which comprises formulating the sulfonylurea compound or its salt by means of various additives, diluting it together with the alkoxyated glyceride with water, and applying it to undesired plants or to a place where they grow.

(15) The above method for controlling undesired plants or inhibiting their growth, which comprises formulating the sulfonylurea compound or its salt by means of various additives, diluting it together with the alkoxyated glyceride and the coadjuvant with water, and applying it to the undesired plants or to a place where they grow.

(16) The above method for controlling undesired plants or inhibiting their growth, which comprises formulating the sulfonylurea compound or its salt, and the alkoxyated glyceride, together with various additives, diluting it with water and applying it to undesired plants or to a place where they grow.

(17) The above method for controlling undesired plants or inhibiting their growth, which comprises formulating the sulfonylurea compound or its salt, the alkoxyated glyceride and the coadjuvant together with
5 various additives, diluting it with water, and applying it to the undesired plants or to a place where they grow.

(18) The above method for controlling undesired plants or inhibiting their growth, characterized by applying foliar application to the undesired plants.

10 (19) The above method for controlling undesired plants or inhibiting their growth, wherein a herbicidally effective amount of other herbicidal compound(s) are further contained.

(20) The above method for enhancing the herbicidal
15 effect of the sulfonylurea compound or its salt by means of the alkoxyated glyceride in an effective amount to enhance the herbicidal activity.

(21) The above method for enhancing the herbicidal effect of the sulfonylurea compound or its salt by means
20 of the alkoxyated glyceride in an effective amount to enhance the herbicidal activity and the coadjuvant in an effective amount to enhance the herbicidal activity.

(22) The above method for enhancing the herbicidal effect of the herbicidal composition comprising the
25 sulfonylurea compound or its salt, and other herbicidal compound(s) by means of the alkoxyated glyceride in an effective amount to enhance the herbicidal activity and

the coadjuvant in an effective amount to enhance the herbicidal activity.

EXAMPLES

EXAMPLE 1

5 (1) Nicosulfuron (purity: 93.6%): 81.78 parts by weight

 (2) Condensate of sodium alkylnaphthalene sulfonate with formalin (tradename: Supragil MNS/90, manufactured by Rhodia Nicca, Ltd.): 5 parts by weight

10 (3) Sodium dodecylbenzene sulfonate (tradename: Neogen Powder, manufactured by DAI-ICHI KOGYO SEIYAKU CO., LTD.): 13.22 parts by weight

 The above components were uniformly mixed, and water was added thereto, followed by kneading, then extrusion
15 granulation, drying and size adjustment to obtain water-dispersible granules. The water-dispersible granules were diluted together with an alkoxylated glyceride with water and then applied.

EXAMPLE 2

20 [1]

 (1) Sodium dodecylbenzenesulfonate (tradename: Sorpol 5060, manufactured by TOHO Chemical Industry Co., Ltd.): 2 parts by weight

 (2) Polyoxyethylene nonylphenyl ether sulfate
25 (tradename: Sorpol 5073, manufactured by TOHO Chemical Industry Co., Ltd.): 3 parts by weight

 (3) Polyoxyethylene dodecylphenyl ether (tradename:

Noigen EA-33, manufactured by DAI-ICHI KOGYO SEIYAKU CO., LTD.): 1 part by weight

(4) Clay (tradename: hydrated clay, manufactured by TODOROKI SANGYO CO., LTD.): 78 parts by weight

5 (5) White carbon (tradename: Carplex #80, manufactured by Shionogi & Co., Ltd.): 16 parts by weight

The above components were mixed to obtain a blend [A].

[2]

10 (1) Foramsulfuron (purity: 99.6%): 10 parts by weight

(2) Blend [A]: 90 parts by weight

The above components were mixed to obtain a wettable powder. The wettable powder was diluted together with an
15 alkoxyated glyceride with water, and then applied.

EXAMPLE 3

(1) Tritosulfuron (purity: at least 98%): 10 parts by weight

(2) Blend [A] obtained in the above Example 2: 90
20 parts by weight

The above components were mixed to obtain a wettable powder. The wettable powder was diluted together with an alkoxyated glyceride with water, and then applied.

EXAMPLE 4

25 (1) Nicosulfuron (purity: 94.3%): 10.7 parts by weight

(2) Polycarboxylate (tradename: Geroxon T/36,

manufactured by Rhodia Nicca, Ltd.): 3 parts by weight

(3) Supragil MNS/90 (tradename): 4.3 parts by weight

(4) Sodium alkylnaphthalenesulfonate (tradename:
Supragil WP, manufactured by Rhodia Nicca, Ltd.): 2 parts
5 by weight

(5) Carplex #80 (tradename): 33.3 parts by weight

(6) Polyoxyethylene hydrogenated castor oil
(tradename: EMANON CH-25, manufactured by Kao
Corporation): 46.7 parts by weight

10 EMANON CH-25 was adsorbed on Carplex #80, and then
other components were mixed thereto to obtain a wettable
powder.

EXAMPLE 5

(1) Nicosulfuron (purity: 94.3%): 10.7 parts by
15 weight

(2) Supragil MNS/90 (tradename): 5 parts by weight

(3) Neogen Powder (tradename): 14.3 parts by weight

(4) Carplex #80 (tradename): 35 parts by weight

(5) Polyoxyethylene hydrogenated castor oil
20 (tradename: EMANON CH-80, manufactured by Kao
Corporation): 35 parts by weight

Molten EMANON CH-80 was adsorbed on Carplex #80, and
then other components were mixed thereto. Water was
added thereto, followed by kneading, then extrusion
25 granulation, drying and size adjustment to obtain water-
dispersible granules.

EXAMPLE 6

(1) Nicosulfuron (purity: 94.3%): 10.7 parts by weight

(2) Supragil MNS/90 (tradename): 5 parts by weight

5 (3) Neogen Powder (tradename): 12.3 parts by weight

(4) Sodium lignin sulfonate (tradename: New Kalgen WG-4, manufactured by TAKEMOTO OIL & FAT CO., LTD.): 2 parts by weight

(5) Carplex #80 (tradename): 35 parts by weight

10 (6) Polyoxyethylene hydrogenated castor oil
pyroglutamic acid isostearate (tradename: PYROTER CPI-60, manufactured by Nihon Emulsion Co., Ltd.): 35 parts by weight

Molten PYROTER CPI-60 was adsorbed on Carplex #80,
15 and then other components were mixed thereto. Water was added thereto, followed by kneading, then extrusion granulation, drying and size adjustment to obtain water-dispersible granules.

EXAMPLE 7

20 (1) Nicosulfuron (purity: 94.3%): 10.7 parts by weight

(2) Supragil MNS/90 (tradename): 5 parts by weight

(3) Calcium dodecylbenzenesulfonate (tradename: Rhodacal 70, manufactured by Rhodia Nicca, Ltd.): 4 parts
25 by weight

(4) Bentonite (tradename: KUNIGEL V1, manufactured by Kunimine Industries Co., Ltd.): 10.3 parts by weight

(5) Carplex #80 (tradename): 35 parts by weight

(6) Polyoxyethylene glyceryl triisostearate
(tradename: EMALOX GWIS-360, manufactured by Nihon
Emulsion Co., Ltd.): 35 parts by weight

5 Molten EMALOX GWIS-360 was adsorbed on Carplex #80,
and then other components were mixed thereto. Water was
added thereto, followed by kneading, then extrusion
granulation, drying and size adjustment to obtain water-
dispersible granules.

10 EXAMPLE 8

(1) Nicosulfuron (purity: 93.4%): 5.35 parts by
weight

(2) Polyoxyethylenetritylphenyl ether phosphoric
acid ester (tradename: Soprophor 3D33, manufactured by
15 Rhodia Nicca, Ltd.): 5 parts by weight

(3) Polydimethylsiloxane (tradename: Rhodorsil
antifoam 432, manufactured by Rhodia Nicca, Ltd.): 0.1
part by weight

(4) Propylene glycol: 5 parts by weight

20 (5) Water: 54.55 parts by weight

(6) EMANON CH-25 (tradename): 30 parts by weight

The above components were mixed and wet-pulverized
for 5 minutes by a wet-grinding machine, to obtain a
water-based suspension concentrate.

25 EXAMPLE 9

(1) Nicosulfuron (purity: 93.4%): 32.11 parts by
weight

(2) Soprophor 3D33 (tradename): 5 parts by weight

(3) Rhodorsil antifoam 432 (tradename): 0.1 parts by weight

(4) Propylene glycol: 5 parts by weight

5 (5) Water: 37.79 parts by weight

(6) EMANON CH-25 (tradename): 20 parts by weight

The above components were mixed and wet-pulverized for 5 minutes by a wet-grinding machine to obtain a water-based suspension concentrate.

10 EXAMPLE 10

(1) Nicosulfuron (purity: 93.4%): 4.93 parts by weight

(2) A mixture of polyoxyethylene hydrogenated castor oil and dioctyl sulfosuccinate: 10.46 parts by weight

15 (3) Bentonite-alkylamino complex (tradename: New D Orben, manufactured by SHIRAIISHI KOGYO KAISHA, LTD.): 1.05 parts by weight

(4) Corn oil: 62.64 parts by weight

(5) EMANON CH-25 (tradename): 20.92 parts by weight

20 The above components were mixed and wet-pulverized for 15 minutes by a wet-grinding machine to obtain an oil-based suspension concentrate.

EXAMPLE 11

(1) Nicosulfuron (purity: 93.4%): 7.38 parts by weight

(2) A mixture of polyoxyethylene hydrogenated castor oil and dioctyl sulfosuccinate: 10.46 parts by weight

(3) New D Orben (tradename): 1.05 parts by weight

(4) Corn oil: 49.73 parts by weight

(5) EMANON CH-25 (tradename): 31.38 parts by weight

The above components were mixed and wet-pulverized
5 for 15 minutes by a wet-grinding machine to obtain an
oil-based suspension concentrate.

EXAMPLE 12

(1) Nicosulfuron (purity: 93.4%): 4.93 parts by
weight

10 (2) A mixture of polyoxyethylene hydrogenated castor
oil and dioctyl sulfosuccinate: 10.46 parts by weight

(3) New D Orben (tradename): 0.53 parts by weight

(4) Urea: 2.09 parts by weight

(4) Corn oil: 61.07 parts by weight

15 (5) EMANON CH-25 (tradename): 20.92 parts by weight

The above components were mixed and wet-pulverized
for 15 minutes by a wet-grinding machine, to obtain an
oil-based suspension concentrate.

EXAMPLE 13

20 (1) Nicosulfuron (purity: 93.4%): 11.7 parts by
weight

(2) A mixture of polyoxyethylene hydrogenated castor
oil and an alkyl aryl sulfonate: 10 parts by weight

(3) Aromatic hydrocarbon solvent (tradename:
25 Solvesso 150, manufactured by EXXON CHEMICAL): 28.3 parts
by weight

(4) EMANON CH-25 (tradename): 50 parts by weight

The above components were mixed and wet-pulverized for 10 minutes by a wet-grinding machine, to obtain an oil-based suspension concentrate.

EXAMPLE 14

5 (1) Nicosulfuron (purity: 93.4%): 11.7 parts by weight

(2) A mixture of polyoxyethylene hydrogenated castor oil and an alkyl aryl sulfonate: 10 parts by weight

(3) Urea: 3 parts by weight

10 (4) Solvesso 150 (tradename): 25.3 parts by weight

(5) EMANON CH-25 (tradename): 50 parts by weight

The above components were mixed and wet-pulverized for 10 minutes by a wet-grinding machine, to obtain an oil-based suspension concentrate.

15 EXAMPLE 15

[1]

(1) Nicosulfuron (purity: 93.4%): 4.93 parts by weight

20 (2) A mixture of polyoxyethylene hydrogenated castor oil and dioctyl sulfosuccinate: 10.46 parts by weight

(3) New D Orben (tradename): 1.05 parts by weight

(4) Corn oil: 62.64 parts by weight

(5) EMANON CH-25 (tradename): 20.92 parts by weight

The above components were mixed and wet-pulverized
25 for 15 minutes by a wet-grinding machine, to obtain a blend [B].

[2]

(1) A mixture of sodium dioctylsulfosuccinate and sodium benzoate (tradename: New Kalgen EX-70, manufactured by TAKEMOTO OIL & FAT CO., LTD.): 50 parts by weight

5 (2) Corn oil: 50 parts by weight

The above components were mixed at 180°C for 30 minutes and then left to cool to obtain a blend [C].

[3]

(1) Blend [B]: 50 parts by weight

10 (2) Blend [C]: 50 parts by weight

The above components were mixed at 80°C for 5 minutes and then left to cool to obtain a gel formulation.

EXAMPLE 16

(1) Nicosulfuron (purity: 93.4%): 7.38 parts by
15 weight

(2) A mixture of polyoxyethylene hydrogenated castor oil and dioctyl sulfosuccinate: 10.46 parts by weight

(3) Urea: 3.14 parts by weight

(4) Corn oil: 52.87 parts by weight

20 (5) EMANON CH-25 (tradename): 26.15 parts by weight

The above components were mixed and wet-pulverized for 10 minutes by a wet-grinding machine, to obtain an oil-based suspension concentrate.

EXAMPLE 17

25 (1) Nicosulfuron (purity: 93.4%): 7.38 parts by weight

(2) A mixture of polyoxyethylene hydrogenated castor

oil and dioctyl sulfosuccinate: 10.46 parts by weight

(3) Urea: 3.14 parts by weight

(4) Corn oil: 47.64 parts by weight

(5) EMANON CH-25 (tradename): 31.38 parts by weight

5 The above components were mixed and wet-pulverized
for 10 minutes by a wet-grinding machine, to obtain an
oil-based suspension concentrate.

EXAMPLE 18

(1) Nicosulfuron (purity: 93.4%): 8.56 parts by
10 weight

(2) Geropon T/36 (tradename): 3 parts by weight

(3) Supragil WP (tradename): 2 parts by weight

(4) Condensate of sodium alkyl naphthalenesulfonate
with formalin (tradename: Supragil MNS/25, Rhodia Nicca,
15 Ltd.): 5 parts by weight

(5) Clay (tradename: MS clay, manufactured by
Fubasamiclay Co., Ltd.): 9.44 parts by weight

(6) EMANON CH-25 (tradename): 40 parts by weight

(7) Carplex #80 (tradename): 32 parts by weight

20 Molten EMANON CH-25 was adsorbed on Carplex #80, and
then other components were mixed thereto to obtain a
wetttable powder.

EXAMPLE 19

(1) Nicosulfuron (purity: 93.4%): 8.56 parts by
25 weight

(2) Geropon T/36 (tradename): 3 parts by weight

(3) Supragil WP (tradename): 2 parts by weight

(4) Supragil MNS/25 (tradename): 5 parts by weight

(5) MS Clay (tradename): 9.44 parts by weight

(6) EMANON CH-80 (tradename): 40 parts by weight

(7) Carplex #80 (tradename): 32 parts by weight

5 Molten EMANON CH-25 was adsorbed on Carplex #80, and then other components were mixed thereto to obtain a wettable powder.

TEST EXAMPLE 1

Upland field soil was filled in a 1/1,000,000 ha
10 pot, seeds of crabgrass (Digitaria sanguinalis L.) were sown therein and grown in a green house. When crabgrass reached 3 leaf stage, a prescribed amount (20 g a.i./ha) of water-dispersible granules containing nicosulfuron as an active ingredient, formulated in accordance with
15 Example 1, was diluted with water corresponding to 300 liters/ha, and an alkoxylated glyceride was added thereto in a concentration of 0.05 wt%, followed by foliar application. Further, for the purpose of comparison, foliar application was carried out in the same manner by
20 using an ethoxylated tallow amine type adjuvant (tradename: Frigate, manufactured by ISK Biosciences Europe S.A.) instead of the alkoxylated glyceride.

On the 21st day after the application of the herbicide, the growth of crabgrass was visually observed
25 (growth inhibition rate (%)=0:untreated plot to 100:complete kill), whereby the results as shown in Table 1 were obtained.

The following is evident from Table 1. Frigate as a commercially available adjuvant enhanced the herbicidal effect of nicosulfuron. Whereas, the alkoxyated glyceride enhanced the herbicidal effect of nicosulfuron more distinctly even at the same concentration.

Table 1

	Alkoxyated glyceride (tradename)	Growth inhibition rate (%)
Present invention	EMANON CH-25	75
	EMANON CH-80	74
	EMALEX GWIS-340	86
	EMALEX GWIS-360	80
	EMALEX GWIS-115	57
	EMALEX GWIS-125	64
	PYROTER GPI-25	59
	PYROTER CPI-60	76
Comparison	Frigate	37
	Nil	3

TEST EXAMPLE 2

Upland field soil was filled in a 1/1,000,000 ha pot, and seeds of black nightshade (Solanum nigrum L.) were sown therein and grown in a green house. When black nightshade reached 2.2 leaf stage, a prescribed amount (20 g a.i./ha) of water-dispersible granules containing nicosulfuron as an active ingredient, formulated in accordance with Example 1, was diluted with water

corresponding to 300 liters/ha, and an alkoxyated glyceride was added thereto in a concentration of 0.1 wt%, followed by foliar application. Further, for the purpose of comparison, foliar application was carried out
5 in the same manner by using a polyoxyethylene sorbitan monolaurate type adjuvant (corresponding to commercially available adjuvant Tween 20, manufactured by NACALAI TESQUE) instead of the alkoxyated glyceride.

On the 21st day after the application of the
10 herbicide, the growth of black nightshade was observed in the same manner as the above Test Example 1, whereby the results as shown in Table 2 were obtained.

The following is evident from Table 2. The polyoxyethylene sorbitan monolaurate as a commercially
15 available adjuvant enhanced the herbicidal effect of nicosulfuron. Whereas, the alkoxyated glyceride enhanced the herbicidal effect of nicosulfuron more distinctly even at the same concentration.

Table 2

	Alkoxyated glyceride (trade name)	Growth inhibition rate (%)
Present invention	EMANON CH-25	57
	EMANON CH-80	83
	EMALEX GWIS-340	69
	EMALEX GWIS-360	81
	EMALEX GWIS-115	42
	EMALEX GWIS-125	60
	PYROTER GPI-25	53
	PYROTER CPI-60	72
Comparison	Polyoxyethylene sorbitan monolaurate	48
	Nil	0

TEST EXAMPLE 3

Upland field soil was filled in a 1/1,000,000 ha
5 pot, and seeds of redroot pigweed (Amaranthus retroflexus
L.) were sown therein and grown in a green house. When
redroot pigweed reached 3.6 leaf stage, a prescribed
amount (5 g a.i./ha) of water-dispersible granules
containing nicosulfuron as an active ingredient,
10 formulated in accordance with the above Example 1, was
diluted with water corresponding to 300 liters/ha, and an
alkoxyated glyceride was added thereto in a
concentration of 0.025 wt%, followed by foliar
application. Further, for the purpose of comparison,

foliar application was carried out in the same manner by using an alkylaryl polyglycol ether type adjuvant (tradename: Citowett, manufactured by BASF France) instead of the alkoxyated glyceride.

5 On the 21st day after the application of the herbicide, the growth of redroot pigweed was observed in the same manner as in Test Example 1, whereby the results as shown in Table 3 were obtained.

The following is evident from Table 3. Citowett as a
10 commercially available adjuvant enhanced the herbicidal effect of nicosulfuron. Whereas, the alkoxyated glyceride enhanced the herbicidal effect of nicosulfuron more distinctly even at the same concentration.

15 Table 3

	Alkoxyated glyceride (tradename)	Growth inhibition rate (%)
Present invention	EMANON CH-25	81
	EMANON CH-80	84
	EMALEX GWIS-340	86
	EMALEX GWIS-360	84
	EMALEX GWIS-115	85
	EMALEX GWIS-125	85
	PYROTER GPI-25	81
	PYROTER CPI-60	86
Comparison	Citowett	76
	Nil	66

TEST EXAMPLE 4

Upland field soil was filled in a 1/1,000,000 ha pot, and seeds of crabgrass were sown therein and grown in a green house. When crabgrass reached 3 leaf stage, a prescribed amount (15 g a.i./ha) of water-dispersible granules containing flazasulfuron as an active ingredient (tradename: KATANA, manufactured by Ishihara Sangyo Kaisha, Ltd.), was diluted with water corresponding to 300 liters/ha, and an alkoxyated glyceride was added thereto in a concentration of 0.025 wt%, followed by foliar application. Further, for the purpose of comparison, foliar application was carried out in the same manner by using Citowett (the same as in Test Example 3) instead of the alkoxyated glyceride.

On the 21st day after the application of the herbicide, the growth of crabgrass was observed in the same manner as in the above Test Example 1, whereby the results as shown in Table 4 were obtained.

The following is evident from Table 4. Citowett as a commercially available adjuvant enhanced the herbicidal effect of flazasulfuron. Whereas, the alkoxyated glyceride enhanced the herbicidal effect of flazasulfuron more distinctly even at the same concentration.

Table 4

	Alkoxyated glyceride (tradename)	Growth inhibition rate (%)
Present invention	EMANON CH-80	89
	EMALEX GWIS-360	84
Comparison	Citowett	68
	Nil	43

TEST EXAMPLE 5

Upland field soil was filled in a 1/1,000,000 ha
5 pot, and seeds of crabgrass were sown therein and grown
in a green house. When crabgrass reached 3 leaf stage, a
prescribed amount (15 g a.i./ha) of water-dispersible
granules containing trifluoxysulfuron as an active
ingredient (tradename: Envoke, manufactured by Syngenta),
10 was diluted with water corresponding to 300 liters/ha,
and an alkoxyated glyceride was added thereto in a
concentration of 0.025 wt%, followed by foliar
application. Further, for the purpose of comparison,
foliar application was carried out in the same manner by
15 using Citowett (the same as in Test Example 3) instead of
the alkoxyated glyceride.

On the 21st day after the application of the
herbicide, the growth of crabgrass was observed in the
same manner as in the above Test Example 1, whereby the
20 results as shown in Table 5 were obtained.

The following is evident from Table 5. Citowett as a

commercially available adjuvant enhanced the herbicidal effect of trifloxysulfuron. Whereas, the alkoxyated glyceride enhanced the herbicidal effect of trifloxysulfuron more distinctly even at the same concentration.

Table 5

	Alkoxyated glyceride (tradename)	Growth inhibition rate (%)
Present invention	EMANON CH-80	89
	EMALEX GWIS-360	90
Comparison	Citowett	82
	Nil	52

TEST EXAMPLE 6

Upland field soil was filled in a 1/1,000,000 ha pot, and seeds of crabgrass were sown therein and grown in a green house. When crabgrass reached 3 leaf stage, a prescribed amount (20 g a.i./ha) of a wettable powder containing foramsulfuron as an active ingredient, formulated in accordance with the above Example 2, was diluted with water corresponding to 300 liters/ha, and the alkoxyated glyceride was added thereto in a concentration of 0.05 wt%, followed by foliar application. Further, for the purpose of comparison, foliar application was carried out in the same manner by using Citowett (the same as in Test Example 3) instead of

the alkoxyated glyceride.

On the 21st day after the application of the herbicide, the growth of crabgrass was observed in the same manner as in the above Test Example 1, whereby the results as shown in Table 6 were obtained.

The following is evident from Table 6. Citowett as a commercially available adjuvant enhanced the herbicidal effect of foramsulfuron. Whereas, the alkoxyated glyceride enhanced the herbicidal effect of foramsulfuron more distinctly even at the same concentration.

Table 6

	Alkoxyated glyceride (tradename)	Growth inhibition rate (%)
Present invention	EMANON CH-80	70
	EMALEX GWIS-360	53
Comparison	Citowett	19
	Nil	3

TEST EXAMPLE 7

Upland field soil was filled in a 1/1,000,000 ha pot, and seeds of velvetleaf (Abutilon theophrasti MEDIC.) were sown therein and grown in a green house. When velvetleaf reached 3 leaf stage, a prescribed amount (20 g a.i./ha) of a wettable powder containing tritosulfuron as an active ingredient, formulated in accordance with the above Example 3, was diluted with

water corresponding to 300 liters/ha, and an alkoxyated glyceride was added thereto in a concentration of 0.05 wt%, followed by foliar application.

On the 21st day after the application of the herbicide, the growth of velvetleaf was observed in the same manner as in the above Test Example 1, whereby the results as shown in Table 7 were obtained.

The following is evident from Table 7. The alkoxyated glyceride enhanced the herbicidal effect of the tritosulfuron distinctly in the same manner as in the preceding Test Examples.

Table 7

	Alkoxyated glyceride (tradename)	Growth inhibition rate (%)
Present invention	EMANON CH-80	74
	EMALEX GWIS-360	66
Comparison	Nil	50

15 TEST EXAMPLE 8

Upland field soil was filled in a 1/1,000,000 ha pot, and seeds of crabgrass were sown therein and grown in a green house. When crabgrass reached 3 leaf stage, a prescribed amount (5 g a.i./ha) of a water-dispersible granules containing rimsulfuron as an active ingredient (tradename: TITUS, manufactured by Du Pont) was diluted with water corresponding to 300 liters/ha, and an

alkoxylated glyceride was added thereto in a concentration of 0.05 wt%, followed by foliar application. Further, for the purpose of comparison, foliar application was carried out in the same manner by using Citowett (the same as in Test Example 3) instead of the alkoxylated glyceride.

On the 21st day after the application of the herbicide, the growth of crabgrass was observed in the same manner as in the above Test Example 1, whereby the results as shown in Table 8 were obtained.

The following is evident from Table 8. Citowett as a commercially available adjuvant enhanced the herbicidal effect of rimsulfuron. Whereas, the alkoxylated glyceride enhanced the herbicidal effect of rimsulfuron more distinctly even at the same concentration.

Table 8

	Alkoxylated glyceride (tradename)	Growth inhibition rate (%)
Present invention	EMANON CH-80	80
	EMALEX GWIS-360	81
Comparison	Citowett	66
	Nil	17

TEST EXAMPLE 9

Upland field soil was filled in a 1/1,000,000 ha pot, and seeds of crabgrass were sown therein and grown

in a green house. When crabgrass reached 3.3 leaf stage, a prescribed amount (35 g a.i./ha) of water-dispersible granules containing nicosulfuron as an active ingredient, formulated in accordance with the above Example 1, was
5 diluted with water corresponding to 300 liters/ha, and an alkoxylated glyceride was added thereto in a concentration of 0.01 wt%, followed by foliar application. Further, for the purpose of comparison, foliar application was carried out in the same manner by
10 using Frigate (the same as in Test Example 1) instead of the alkoxylated glyceride.

On the 22nd day after the application of the herbicide, the growth of crabgrass was observed in the same manner as in the above Test Example 1, whereby the
15 results as shown in Table 9 were obtained.

The following is evident from Table 9. Frigate as a commercially available adjuvant enhanced the herbicidal effect of nicosulfuron. Whereas, the alkoxylated glyceride enhanced the herbicidal effect of nicosulfuron
20 more distinctly even at the same concentration.

Table 9

	Alkoxyated glyceride (tradename)	Growth inhibition rate (%)
Present invention	EMANON CH-25	78
	EMANON CH-80	78
	EMALEX GWIS-360	82
	EMALEX TPIS-350	66
Comparison	Frigate	35
	Nil	23

TEST EXAMPLE 10

Upland field soil was filled in a 1/1,000,000 ha
 5 pot, and seeds of crabgrass were sown therein and grown
 in a green house. When crabgrass reached 3.3 leaf stage,
 a prescribed amount (30 g a.i./ha) of water-dispersible
 granules containing nicosulfuron as an active ingredient,
 formulated in accordance with the above Example 1, was
 10 diluted with water corresponding to 300 liters/ha, and an
 alkoxyated glyceride was added thereto in a
 concentration of 0.01 wt%, followed by foliar
 application. Further, for the purpose of comparison,
 foliar application was carried out in the same manner by
 15 using Citowett (the same as in Test Example 3) instead of
 the alkoxyated glyceride.

On the 22nd day after the application of the
 herbicide, the growth of crabgrass was observed in the
 same manner as in the above Test Example 1, whereby the

results as shown in Table 10 were obtained.

The following is evident from Table 10. Citowett as a commercially available adjuvant did not enhance the herbicidal effect of nicosulfuron. Whereas, the
5 alkoxyated glyceride enhanced the herbicidal effect of nicosulfuron distinctly even at the same concentration.

Table 10

	Alkoxyated glyceride (tradename)	Growth inhibition rate (%)
Present invention	EMANON CH-25	75
	EMANON CH-80	79
	EMALEX GWIS-360	79
	EMALEX TPIS-350	59
Comparison	Citowett	25
	Nil	25

10 TEST EXAMPLE 11

Upland field soil was filled in a 1/1,000,000 ha pot, and seeds of crabgrass were sown therein and grown in a green house. When crabgrass reached 3.3 leaf stage, a prescribed amount (30 g a.i./ha) of water-dispersible
15 granules containing nicosulfuron as an active ingredient, formulated in accordance with the above Example 1, was diluted with water corresponding to 300 liters/ha, and an alkoxyated glyceride was added thereto in a concentration of 0.02 wt%, followed by foliar

application. Further, for the purpose of comparison, foliar application was carried out in the same manner by using Citowett (the same as in Test Example 3) instead of the alkoxylated glyceride.

5 On the 22nd day after the application of the herbicide, the growth of crabgrass was observed in the same manner as in the above Test Example 1, whereby the results as shown in Table 11 were obtained.

The following is evident from Table 11. Citowett as
10 a commercially available adjuvant did not enhance the herbicidal effect of nicosulfuron. Whereas, the alkoxylated glyceride enhanced the herbicidal effect of nicosulfuron distinctly even at the same concentration.

15 Table 11

	Alkoxylated glyceride (tradename)	Growth inhibition rate (%)
Present invention	EMANON CH-25	89
	EMANON CH-80	87
	EMALEX GWIS-360	88
	EMALEX TPIS-350	84
Comparison	Citowett	25
	Nil	25

TEST EXAMPLE 12

Upland field soil was filled in a 1/1,000,000 ha pot, and seeds of barnyard grass (Echinochloa crus-galli

L.) were sown therein and grown in a green house. When barnyard grass reached 2 leaf stage, a prescribed amount (1.5 g a.i./ha) of water-dispersible granules containing nicosulfuron as an active ingredient, formulated in accordance with the above Example 1, was diluted with water corresponding to 300 liters/ha, and an alkoxyated glyceride was added thereto in a concentration of 0.3 wt%, followed by foliar application. Further, for the purpose of comparison, foliar application was carried out in the same manner by using Citowett (the same as in Test Example 3) instead of the alkoxyated glyceride.

On the 22nd day after the application of the herbicide, the growth of barnyard grass was observed in the same manner as in the above Test Example 1, whereby the results as shown in Table 12 were obtained.

The following is evident from Table 12. Citowett as a commercially available adjuvant enhanced the herbicidal effect of nicosulfuron. Whereas, the alkoxyated glyceride enhanced the herbicidal effect of nicosulfuron more distinctly even at the same concentration.

Table 12

	Alkoxyated glyceride (tradename)	Growth inhibition rate (%)
Present invention	EMANON CH-25	94
	EMANON CH-80	89
	EMALEX GWIS-360	96
	EMALEX TPIS-350	98
Comparison	Citowett	56
	Nil	30

TEST EXAMPLE 13

Upland field soil was filled in a 1/1,000,000 ha
 5 pot, and seeds of barnyard grass were sown therein and
 grown in a green house. When barnyard grass reached 2
 leaf stage, a prescribed amount (1.5 g a.i./ha) of water-
 dispersible granules containing nicosulfuron as an active
 ingredient, formulated in accordance with the above
 10 Example 1, was diluted with water corresponding to 300
 liters/ha, and an alkoxyated glyceride was added thereto
 in a concentration of 0.2 wt%, followed by foliar
 application. Further, for the purpose of comparison,
 foliar application was carried out in the same manner by
 15 using Citowett (the same as in Test Example 3) instead of
 the alkoxyated glyceride.

On the 22nd day after the application of the
 herbicide, the growth of barnyard grass was observed in
 the same manner as in the above Test Example 1, whereby

the results as shown in Table 13 were obtained.

The following is evident from Table 13. Citowett as a commercially available adjuvant enhanced the herbicidal effect of nicosulfuron. Whereas, the alkoxy-
5 glyceride enhanced the herbicidal effect of nicosulfuron more distinctly even at the same concentration.

Table 13

	Alkoxy- glyceride (tradename)	Growth inhibition rate (%)
Present invention	EMANON CH-25	93
	EMANON CH-80	86
	EMALEX GWIS-360	96
	EMALEX TPIS-350	98
Comparison	Citowett	35
	Nil	30

10 TEST EXAMPLE 14

Upland field soil was filled in a 1/1,000,000 ha
pot, and seeds of barnyard grass were sown therein and
grown in a green house. When barnyard grass reached 2
leaf stage, a prescribed amount (1.5 g a.i./ha) of water-
15 dispersible granules containing nicosulfuron as an active
ingredient, formulated in accordance with the above
Example 1, was diluted with water corresponding to 300
liters/ha, and an alkoxy- glyceride was added thereto
in a concentration of 0.1 wt%, followed by foliar

application. Further, for the purpose of comparison, foliar application was carried out in the same manner by using Frigate (the same as in Test Example 1) instead of the alkoxyated glyceride.

5 On the 22nd day after the application of the herbicide, the growth of barnyard grass was observed in the same manner as in the above Test Example 1, whereby the results as shown in Table 14 were obtained.

The following is evident from Table 14. Frigate as a
10 commercially available adjuvant enhanced the herbicidal effect of nicosulfuron. Whereas, the alkoxyated glyceride enhanced the herbicidal effect of nicosulfuron more distinctly even at the same concentration.

15 Table 14

	Alkoxyated glyceride (tradename)	Growth inhibition rate (%)
Present invention	EMANON CH-25	91
	EMANON CH-80	93
	EMALEX GWIS-360	95
	EMALEX TPIS-350	95
Comparison	Frigate	53
	Nil	30

TEST EXAMPLE 15

Upland field soil was filled in a 1/1,000,000 ha pot, and seeds of barnyard grass were sown therein and

grown in a green house. When barnyard grass reached 2 leaf stage, a prescribed amount (3 g a.i./ha) of water-dispersible granules containing nicosulfuron as an active ingredient, formulated in accordance with the above

5 Example 1, was diluted with water corresponding to 300 liters/ha, and an alkoxyated glyceride was added thereto in a concentration of 0.1 wt%, followed by foliar application. Further, for the purpose of comparison, foliar application was carried out in the same manner by

10 using Citowett (the same as in Test Example 3) instead of the alkoxyated glyceride.

On the 22nd day after the application of the herbicide, the growth of barnyard grass was observed in the same manner as in the above Test Example 1, whereby

15 the results as shown in Table 15 were obtained.

The following is evident from Table 15. Citowett as a commercially available adjuvant enhanced the herbicidal effect of nicosulfuron. Whereas, the alkoxyated glyceride enhanced the herbicidal effect of nicosulfuron

20 more distinctly even at the same concentration.

Table 15

	Alkoxyated glyceride (tradename)	Growth inhibition rate (%)
Present invention	EMANON CH-25	94
	EMANON CH-80	99
	EMALEX GWIS-360	97
	EMALEX TPIS-350	97
Comparison	Citowett	42
	Nil	28

TEST EXAMPLE 16

Upland field soil was filled in a 1/1,000,000 ha
 5 pot, and seeds of crabgrass were sown therein and grown
 in a green house. When crabgrass reached 3 leaf stage, a
 prescribed amount (18 g a.i./ha) of water-dispersible
 granules containing nicosulfuron as an active ingredient,
 formulated in accordance with the above Example 1, was
 10 diluted with water corresponding to 300 liters/ha, and an
 alkoxyated glyceride and a chelating agent (citric acid
 or ethylene diamine tetraacetic acid (EDTA)) as a
 coadjuvant were added thereto in prescribed
 concentrations, followed by foliar application.

15 On the 22nd day after the application of the
 herbicide, the growth of crabgrass was observed in the
 same manner as in the above Test Example 1, whereby the
 results as shown in Table 16 were obtained.

The following is evident from Table 16. The

alkoxylated glyceride enhanced the herbicidal effect of nicosulfuron. However, when the coadjuvant was further added, the herbicidal effect of nicosulfuron was enhanced more distinctly in spite of the fact that the total

5 amount of the alkoxylated glyceride and the coadjuvant was the same as the amount of the alkoxylated glyceride added alone.

Table 16

Alkoxylated glyceride (tradename)	Chelating agent	Concentration in diluted liquid (wt%)		Growth inhibition rate (%)
		Alkoxylated glyceride	Chelating agent	
EMANON CH-80	-	0.02	-	75
	-	0.025	-	82
	Citric acid	0.015	0.005	75
	Citric acid	0.02	0.005	93
	EDTA	0.015	0.005	80
	EDTA	0.02	0.005	94

10

TEST EXAMPLE 17

Upland field soil was filled in a 1/1,000,000 ha pot, and seeds of crabgrass were sown therein and grown in a green house. When crabgrass reached 3 leaf stage, a
 15 prescribed amount (20 g a.i./ha) of water-dispersible

granules containing nicosulfuron as an active ingredient, formulated in accordance with the above Example 1, was diluted with water corresponding to 300 liters/ha, and an alkoxyated glyceride was added thereto in a

5 concentration of 0.025 wt%, followed by foliar application. Further, for the purpose of comparison, foliar application was carried out in the same manner by using Citowett (the same as in Test Example 3) instead of the alkoxyated glyceride.

10 On the 21st day after the application of the herbicide, the growth of crabgrass was observed in the same manner as in the above Test Example 1, whereby the results as shown in Table 17 were obtained.

The following is evident from Table 17. Citowett as
15 a commercially available adjuvant enhanced the herbicidal effect of nicosulfuron. Whereas, the alkoxyated glyceride enhanced the herbicidal effect of nicosulfuron more distinctly even at the same concentration.

Table 17

	Alkoxyated glyceride (tradename)	Growth inhibition rate (%)
Present invention	NIKKOL HCO-5	85
	Sorpol HC-10	89
	Sorpol HC-20	91
	PEGNOL HC-30	92
	Sorpol HC-40	94
	Sorpol HC-50	93
	Sorpol HC-80	94
	Sorpol HC-100	93
	Sorpol HC-150	93
Comparison	Citowett	8
	Nil	2

TEST EXAMPLE 18

Upland field soil was filled in a 1/1,000,000 ha. pot, and seeds of crabgrass were sown therein and grown in a green house. When crabgrass reached 3 leaf stage, a prescribed amount (15 g a.i./ha) of the one obtained by the formulation method in the above Example 17, was diluted with water corresponding to 300 liters/ha, followed by foliar application. Further, for the purpose of comparison, foliar application was carried out in the same manner using a prescribed amount (15 g a.i./ha) of Onehope NYUZAI (manufactured by ISK BIOSCIENCES K.K.) as a commercial product containing nicosulfuron as an active ingredient.

On the 21st day after the application of the herbicide, the growth of crabgrass was observed in the same manner as in the above Test Example 1, whereby the results as shown in Table 18 were obtained.

- 5 The following is evident from Table 18. The oil-based suspension concentrate of the present invention containing the alkoxyated glyceride exhibited a distinctive herbicidal effect, even at a dose (15 g a.i./ha) lower than the standard dose of Onehope NYUZAI.

10

Table 18

	Formulation	Growth inhibition rate (%)
Present invention	Example 17	72
Comparison	Onehope NYUZAI	48

TEST EXAMPLE 19

- Upland field soil was filled in a 1/1,000,000 ha pot, and seeds of crabgrass were sown therein and grown in a green house. When crabgrass reached 3 leaf stage, a prescribed amount (20 g a.i./ha) of water-dispersible granules containing nicosulfuron as an active ingredient, formulated in accordance with the above Example 1, was diluted with water corresponding to 300 liters/ha, and an alkoxyated glyceride and a nitrogen-containing fertilizer (tradename: ANBLA, manufactured by Sankyo Co., Ltd., ammoniacal nitrogen:water-soluble
- 15
- 20

phosphoric acid:water-soluble potassium = 4:30:16)
 were added thereto at prescribed concentrations,
 followed by foliar application.

On the 21st day after the application of the
 5 herbicide, the growth of crabgrass was observed in the
 same manner as in the above Test Example 1, whereby the
 results as shown in Table 19 were obtained.

The following is evident from Table 19. The
 alkoxyated glyceride enhanced the herbicidal effect of
 10 nicosulfuron. However, when the nitrogen-containing
 fertilizer was added, the herbicidal effect of
 nicosulfuron was further enhanced.

Table 19

Alkoxyated glyceride (tradename)	Nitrogen- containing fertilizer (tradename)	Concentration in diluted liquid (wt%)		Growth inhibition rate (%)
		Alkoxyated glyceride	Nitrogen- containing fertilizer	
EMANON	-	0.025	-	90
CH-80	ANBLA	0.025	0.005	93